

Last day

* Complex exponentials

Let $z = x + iy$,

$$e^{iz} = e^{i(x+iy)} = e^{ix-y} = e^{-y} e^{ix}$$

$$= e^{-y} (\cos(x) + i \sin(x))$$

$$= e^{-y} \cos(x) + i e^{-y} \sin(x)$$

If $x = \operatorname{Re}(z) = 0$, $z = iy$

$$e^z = e^{iy} = \cos(y) + i \sin(y).$$


* polar representation of complex numbers

Let $z = x + iy$, then in polar coordinates

$$z = r e^{i\theta}$$

where $r = |z|$

θ (argument of z) is the angle z makes with the positive x -axis.



EIGENVALUES AND EIGENVECTORS


Let A be an $n \times n$ matrix. A number λ and a vector \vec{x} is an eigenvalue eigenvector pair of A if;

(i) $\vec{x} \neq \vec{0}$

(ii) $A\vec{x} = \lambda\vec{x}$

If \exists (there exist) such λ and a vector \vec{x} , then λ is an eigenvalue of A with corresponding eigenvector \vec{x} .


Consider the reflection matrix, R , about a line that ~~ma~~ makes an angle of θ with the x -axis, and let \vec{x} be a vector on the line



$$R\vec{x} = \vec{x} \quad \text{---} \quad (*)$$

relate this to $A\vec{x} = \lambda\vec{x}$.

This shows that $(*)$ is an eigenvalue problem where $\lambda = 1$ is an eigenvalue with corresponding eigen vector \vec{x} .



$$R\vec{y} = -\vec{y}$$

$\Rightarrow \lambda = -1$ is an eigenvalue with eigen vector \vec{y} .

We observe that these are the only two vectors that can give us a scalar multiple which when multiplied by A will give us a scalar multiple of the original vector.

This implies that the reflection matrix has only two eigen values $\lambda_1 = 1$, $\lambda_2 = -1$ with corresponding eigen vectors \vec{x} and \vec{y} , respectively.

These eigenvectors are linearly independent and form a basis for \mathbb{R}^2 .

They also form a basis of eigenvectors of the reflection matrix.

Let A be an $n \times n$ matrix and λ be an eigenvalue of A with eigenvector \vec{x} . Then

$$A\vec{x} = \lambda\vec{x}$$

Let ~~α~~ $\alpha \neq 0$ be a scalar.

$$A(\alpha\vec{x}) = A\alpha\vec{x} = \alpha A\vec{x} = \alpha\lambda\vec{x}$$

$$A(\alpha\vec{x}) = \lambda(\alpha\vec{x})$$


$\therefore \alpha\vec{x}$ is also an eigenvector of eigenvalue λ .

This implies that the eigenvector of an eigenvalue is not uniquely determined.

And this also shows that the direction of the eigenvector is the important factor.

Let us consider the rotation matrix. Let R be the matrix of rotation by an angle 45° .

Let \vec{x} be a vector



Observe that the direction of \vec{x} is changed by R .

$\Rightarrow R$ does not have an eigenvalue.

COMPUTING the eigenvalue and eigenvectors

Let A be an $n \times n$ matrix with eigenvalue λ and corresponding eigenvector \vec{x} .

Then
$$A\vec{x} = \lambda\vec{x}$$

$$A\vec{x} - \lambda\vec{x} = \vec{0}$$

$$\text{or } (A - \lambda I)\vec{x} = \vec{0}$$

Let $B = A - \lambda I$, then

$$B\vec{x} = \vec{0} \quad (\text{homogeneous system})$$

If $\vec{x} \neq \vec{0}$, then $\det(B) = 0$

$$\Rightarrow \det(A - \lambda I) = 0$$

I_n is identity matrix.

this is called the characteristic equation.

After we obtain the eigenvalue, we can

use $(A - \lambda I) \vec{x} = \vec{0}$

to get the eigenvectors.

Example: Find the eigenvalues and eigenvectors of the matrix $A = \begin{pmatrix} 2 & 3 \\ 3 & 2 \end{pmatrix}$

Solution:

Let λ be an eigenvalue of A .

then $(A - \lambda I) \vec{x} = \vec{0}$

$$|A - \lambda I| = 0$$

$$\left| \begin{pmatrix} 2 & 3 \\ 3 & 2 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right| = 0$$

$$\begin{vmatrix} 2-\lambda & 3 \\ 3 & 2-\lambda \end{vmatrix} = 0$$

$$(2-\lambda)(2-\lambda) - 9 = 0$$

$$\lambda^2 - 4\lambda - 5 = 0$$

$$(\lambda - 5)(\lambda + 1) = 0$$

$$\lambda_1 = 5 \quad \text{and} \quad \lambda_2 = -1$$

\therefore these are the eigenvalues of A.

Let us find the eigenvector.

$$\text{For } \lambda_1 = 5, \quad (A - \lambda_1 I) \vec{x} = \vec{0}$$

$$\left(\begin{pmatrix} 2 & 3 \\ 3 & 2 \end{pmatrix} - 5 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right) \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} -3 & 3 \\ 3 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\left(\begin{array}{cc|c} -3 & 3 & 0 \\ 3 & -3 & 0 \end{array} \right) \sim \left(\begin{array}{cc|c} 3 & -3 & 0 \\ 3 & -3 & 0 \end{array} \right)$$

$$\sim \left(\begin{array}{cc|c} 3 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right)$$

$$\Rightarrow 3x_1 - 3x_2 = 0$$

$$\Rightarrow x_1 = x_2$$

$$\vec{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_2 \end{pmatrix} = x_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}, x_2 \text{ is free}$$

$$\Rightarrow \vec{x} = \alpha \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (\alpha \neq 0, \text{ is constant})$$

$$\lambda_2 = -1, \quad (A - \lambda_2 I) \vec{x} = \vec{0}$$

$$\left(\begin{pmatrix} 2 & 3 \\ 3 & 2 \end{pmatrix} - (-1) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right) \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 3 & 3 \\ 3 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\left(\begin{array}{cc|c} 3 & 3 & 0 \\ 3 & 3 & 0 \end{array} \right) \sim \left(\begin{array}{cc|c} 3 & 3 & 0 \\ 0 & 0 & 0 \end{array} \right)$$

$$3x_1 + 3x_2 = 0$$

$$x_1 = -x_2$$

$$\vec{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} -x_2 \\ x_2 \end{pmatrix} = x_2 \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

$$\vec{x} = \beta \begin{pmatrix} -1 \\ 1 \end{pmatrix}, \quad (\beta \neq 0, \text{ constant})$$

We have two eigenvalues $\lambda_1 = 5$ and $\lambda_2 = -1$
with corresponding eigenvectors

$$\vec{x}_1 = \alpha \begin{pmatrix} 1 \\ 1 \end{pmatrix} \text{ and } \vec{x}_2 = \beta \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

respectively.

$\alpha \neq 0$ and $\beta \neq 0$ are scalars.

Example : (Repeated eigenvalues)

Find the eigenvalues and eigenvectors of

$$A = \begin{pmatrix} 0 & -4 & -6 \\ 4 & 0 & -3 \\ 1 & 2 & 5 \end{pmatrix}$$

Let λ be an eigenvalue of A , then

$$|A - \lambda I| = 0$$

$$\Rightarrow \left| \begin{pmatrix} 0 & -4 & -6 \\ 4 & 0 & -3 \\ 1 & 2 & 5 \end{pmatrix} \right| \rightarrow \left| \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right| = 0$$

$$\Rightarrow \begin{vmatrix} -\lambda & -4 & -6 \\ -4 & -\lambda & -3 \\ 1 & 2 & 5-\lambda \end{vmatrix} = 0$$

$$-\lambda[-\lambda(5-\lambda)+6] + 4[-1(5-\lambda)+3] - 6[-2+\lambda] = 0$$

Simplifying, we get

$$\lambda^3 - 5\lambda^2 + 8\lambda - 4 = 0$$

(characteristic polynomial)

By ~~map~~ inspection, we get $\lambda = 1$ as a root of the polynomial. $\Rightarrow \lambda - 1 = 0$

$$\begin{array}{r} \lambda^2 - 4\lambda + 4 \\ \lambda - 1 \overline{) \lambda^3 - 5\lambda^2 + 8\lambda - 4} \\ \underline{-(\lambda^3 - \lambda^2)} \\ 0 - 4\lambda^2 + 8\lambda - 4 \\ \underline{-(-4\lambda^2 + 4\lambda)} \\ 4\lambda - 4 \\ \underline{4\lambda - 4} \\ 0 \quad 0 \end{array}$$

$$\Rightarrow (\lambda - 1)(\lambda^2 - 4\lambda + 4) = 0$$

$$(\lambda - 1)(\lambda - 2)(\lambda - 2) = 0$$

$$(\lambda - 1)(\lambda - 2)^2 = 0$$

\therefore the eigenvalues are

$$\lambda_1 = 1 \quad \text{and} \quad \lambda_2 = 2$$

$$\text{For } \lambda_1 = 1, \quad (A - \lambda_1 I) \vec{x} = \vec{0}$$

$$\Rightarrow \left[\begin{pmatrix} 0 & -4 & -6 \\ -1 & 0 & -3 \\ 1 & 2 & 5 \end{pmatrix} - I \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right] \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} -1 & -4 & -6 \\ -1 & -1 & -3 \\ 1 & 2 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\left[\begin{array}{ccc|c} -1 & -4 & -6 & 0 \\ -1 & -1 & -3 & 0 \\ 1 & 2 & 4 & 0 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 2 & 4 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

from row 2, $x_2 + x_3 = 0 \Rightarrow x_2 = -x_3$
from row 1

$$x_1 + 2x_2 + 4x_3 = 0$$

$$x_1 = -2x_2 - 4x_3 = -2(-x_3) - 4x_3 = 2x_3 - 4x_3$$

$$x_1 = -2x_3$$

$$\vec{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -2x_3 \\ -x_3 \\ x_3 \end{pmatrix} = x_3 \begin{pmatrix} -2 \\ -1 \\ 1 \end{pmatrix}$$

x_3 is free.

$$\text{For } \lambda_2 = 2, \quad (A - \lambda_2 I) \vec{X} = \vec{0}$$

$$\Rightarrow \left(\begin{pmatrix} 0 & -4 & -6 \\ -1 & 0 & -3 \\ 1 & 2 & 5 \end{pmatrix} - 2 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\left[\begin{array}{ccc|c} -2 & -4 & -6 & 0 \\ -1 & -2 & -3 & 0 \\ 1 & 2 & 3 & 0 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 1 & 2 & 3 & 0 \\ 1 & 2 & 3 & 0 \end{array} \right] \begin{array}{l} R_1 / -2 \\ +R_2 \\ +R_3 \end{array}$$

$$\sim \left[\begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$x_1 + 2x_2 + 3x_3 = 0$$

$$x_1 = -2x_2 - 3x_3$$

$$\vec{X} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -2x_2 - 3x_3 \\ x_2 \\ x_3 \end{pmatrix} = x_2 \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix}$$

x_2 and x_3 are free.

x_2 and x_3 cannot both be zero. ~~at least~~

∴ the eigenvector of $\lambda_2 = 2$ is

$$\vec{X} = \alpha_1 \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix} + \alpha_2 \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix}$$

where α_1 and α_2 are scalars and cannot both be zero.

For example: take $\alpha_1 = 0$, and $\alpha_2 = 1$,

$$\vec{X} = \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix} \text{ is an eigenvector of } \lambda_2$$

OR you can take $\alpha_1 = 1$ and $\alpha_2 = 0$

$$\vec{X} = \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix} \text{ is also an eigenvector of } \lambda_2.$$